

Oscillations of Bose condensates in a one-dimensional optical superlattice

Chou-Chun Huang and Wen-Chin Wu

Department of Physics, National Taiwan Normal University, Taipei 11650, Taiwan

Oscillations of atomic Bose-Einstein condensates in a 1D optical lattice with a two-point basis is investigated. In the low-frequency regime, four branches of modes are resolved, that correspond to the transverse in-phase and out-of-phase breathing modes, and the longitudinal acoustic and optical phonon modes of the condensates. Dispersions of these modes depend intimately on the values of two intersite Josephson tunneling strengths, J_1 and J_2 , and the on-site repulsion U between the atoms. Observation of these mode dispersions is thus a direct way to access them.

PACS numbers: 03.75.Hh, 32.80.Pj, 03.65.-w

Cold atoms in optical lattices is currently an active research topics in the cross-disciplined field of atomic molecular and optics physics and condensed-matter physics. From the condensed matter point of view, because (i) laser power can be tuned to vary the ratio of on-site repulsion to inter-site coupling of cold atoms, (ii) different geometry of laser beams can be taken to manipulate different dimensions and configurations of optical lattices, (iii) different types of atoms (fermions, bosons, ions, or their mixtures) can be loaded into the optical lattice, consequently the simulation of condensed matter environment is considerably easy and flexible. Recent investigations on the cold-atom system with an optical lattice have been made in various aspects, including the superfluid (SF)-Mott insulator quantum phase transition^{1,2}, the band-structure phenomena^{3,4}, and the quantum informatics^{5,6}.

In this paper, we consider a physical environment very common in crystalline solids. It is often in a solid that the system is best described by a superlattice (*i.e.*, lattice with a basis). This occurs when there are more than one type of atoms (or sites) within a unit cell. Correspondingly in the cold-atom system, a 1D optical lattice with n -point basis can be set up as follows. One can shine two laser beams coherently towards the same direction (say, z -axis), with one having n times of frequency (and thus n times of wave vector) of the other. Along with their reflected beams, the resulting electric field is $E(z, t) = A_1 \cos(nkz)e^{i\omega t} + A_2 \cos(kz)e^{i\omega t} + \text{c.c.}$ and the dipole energy, which is proportional to the time average of electric field square, is thus

$$U(z) \propto \overline{E(z, t)^2} \sim A_1^2 \cos^2(nkz) + A_2^2 \cos^2(kz). \quad (1)$$

The cross term vanishes due to time average. Consequently Bose condensates, which is pre-confined in the x - y plane by the magnetic trap, can be optically trapped and form a z -direction 1D optical lattice with an n -point basis. Since the ratio of amplitude A_1/A_2 can be adjusted and the factor n can be chosen, 1D optical superlattice can be created for a great flexibility. Figure 1 illustrates how the 1D optical superlattice with a 2 and 3-point basis is formed and their resulting potential energies ($A_1/A_2 = \sqrt{5}$ is taken). Josephson coupling J 's that differ due to condensates tunnelling across different

potential barriers are also indicated.




FIG. 1: (a) Illustration of how a 1D optical lattice with an n -point basis is formed by two laser beams and their reflected beams. Frame (b) and (c) show the resulting potential energies (thick solid lines) as a function of lattice spacing of the $n = 2$ and $n = 3$ case. c denotes the lattice constant and J 's denote the Josephson tunnelling couplings. In frame (b), d denotes the spacing between any potential maximum and its nearby minimum.

Since the achievement of atomic BEC in 1995, dynamics of single Bose condensate had been under detailed investigation both theoretically and experimentally⁷. When condensates are distributed in an optical lattice, in addition to collective modes, phonons can also propagate among the lattice. How the collective and phonon modes are coupled in such a condensed system thus has attracted a large number of studies^{3,8,9,10,11,12}. In fact, condensate excitations in a 3D optical lattice have been measured recently in the SF regime¹³. Among different theoretical approaches, Martikainen and Stoof¹² have recently applied the variational approach to study the transverse breathing mode and longitudinal acoustic phonon in a simple 1D Bose-condensed optical lattice. The advantage of this approach is that it enables naturally the coupling between the breathing and phonon modes, and at the same time, allows for analytical results. In the present context, we shall generalize their approach to study how the breathing and phonon modes

of Bose condensates are coupled and behave in a 1D optical superlattice.

At sufficiently low temperatures, dynamics of Bose condensates is governed by the time-dependent Gross-Pitavetskii (GP) energy functional

$$E[\Psi^*, \Psi] = \int d^3r \left\{ -\frac{\hbar^2}{2m} \Psi^* \nabla^2 \Psi + [V_{\text{trap}}(\mathbf{r}) + \frac{U_0}{2} |\Psi|^2 - \mu] |\Psi|^2 \right\}, \quad (2)$$

where $\Psi = \Psi(\mathbf{r}, t)$ is the time-dependent wave function of the condensate, μ is the chemical potential, and $U_0 = 4\pi\hbar^2 a/m$ with a the s -wave scattering length of the two-body interaction. In the present case of interest, the external trap potential has two contributions, $V_{\text{trap}}(\mathbf{r}) = V_m(\mathbf{r}) + V_o(\mathbf{r})$, where $V_m(\mathbf{r}) = \frac{1}{2}m[\omega_\rho^2 \rho^2 + \omega_z^2 z^2]$ with $\rho^2 \equiv x^2 + y^2$ is due to the magnetic trap, while $V_o(\mathbf{r}) = V_o(z) = V_1 \cos^2(\frac{2\pi z}{c}) + V_2 \cos^2(\frac{\pi z}{c})$ is due to the optical trap. When $\omega_\rho \gg \omega_z$ for the magnetic trap, optical potential then results a 1D optical superlattice (along z axis) with a two-point basis. Here c corresponds to the lattice constant which is related to the wavelength (λ) or the wavevector (k) of the laser beam with frequency ω by $c = \lambda/2 = \pi/k$ (see Fig. 1). It is convenient to work with dimensionless quantities and rescale the energy, time, and length as $E/(\hbar\omega_\rho) \rightarrow E$, $t\omega_\rho \rightarrow t$, and $\mathbf{r}/\ell_\rho \rightarrow \mathbf{r}$ ($\ell_\rho \equiv \sqrt{\hbar/m\omega_\rho}$). Eq. (2) thus becomes

$$E[\Psi^*, \Psi] = \int d^3r \left\{ -\frac{1}{2} \Psi^* \nabla^2 \Psi + \left[\frac{1}{2}(x^2 + y^2) + \bar{V}_o(z) + \frac{\bar{U}_0}{2} |\Psi|^2 - \bar{\mu} \right] |\Psi|^2 \right\}, \quad (3)$$

where the dimensionless coupling $\bar{V}_o(z) \equiv V_o(z)/(\hbar\omega_\rho)$, $\bar{U}_0 \equiv 4\pi a/\ell_\rho$, and $\bar{\mu} \equiv \mu/(\hbar\omega_\rho)$. The frequency ω_z is completely dropped in (3).

For longitudinal phonon modes to be considered under the *harmonic* approximation, *tight-binding* (TB) limit is applied. This means that anharmonic effect is negligibly small. This also means that optical potential barriers need to be large, so condensate wave functions are strongly localized in z -direction around the potential minimums. In this limit, $\Psi(\mathbf{r}, t)$ can be taken to be the form

$$\Psi(\mathbf{r}, t) = \sum_{\ell} \{ w_1[z - \ell c - d] \Phi_1(x, y, \ell; t) + w_2[z - (\ell + 1)c + d] \Phi_2(x, y, \ell; t) \}, \quad (4)$$

where ℓ sums over all unit cells and d is the spacing between any potential maximum and its nearby minimum (see Fig. 1). Pertaining to site 1 and site 2, w_1 and w_2 are the strongly localized Wannier functions depending on z only, while Φ_1 and Φ_2 are the ones associated with x, y coordinates which in turn contain the periodic factor $\sim \exp(i\ell k c)$ for any propagating wave (along

the lattice direction) of wave vector k . It is noted that $w_1(z) = w_2(-z)$ for the reflection symmetry. Moreover, for simplicity, fluctuations (time dependence) are considered to be through $\Phi_i(x, y, \ell; t)$ only.

Substitution of (4) reduces (3) into

$$E = \sum_{\ell} \sum_{i=1,2} \int d^2\rho \left\{ -\frac{1}{2} \Phi_i^*(\ell) \nabla^2 \Phi_i(\ell) + \left[\frac{1}{2} \rho^2 + \frac{U}{2} |\Phi_i(\ell)|^2 - \bar{\mu} \right] |\Phi_i(\ell)|^2 \right\} + \sum_{\ell} \int d^2\rho [J_1 \Phi_1^*(\ell) \Phi_2(\ell - 1) + \text{c.c.}] + \sum_{\ell} \int d^2\rho [J_2 \Phi_1^*(\ell) \Phi_2(\ell) + \text{c.c.}], \quad (5)$$

where $\Phi_i(\ell) \equiv \Phi_i(x, y, \ell; t)$ and

$$J_1 \equiv - \int dz w_1^*(z - d) \left[-\frac{1}{2} \frac{\partial^2}{\partial z^2} + \bar{V}_o(z) \right] w_2(z + d) \\ J_2 \equiv - \int dz w_1^*(z - d) \left[-\frac{1}{2} \frac{\partial^2}{\partial z^2} + \bar{V}_o(z) \right] w_2(z - c + d) \\ U \equiv \bar{U}_0 \int dz |w_1(z)|^4 = \bar{U}_0 \int dz |w_2(z)|^4. \quad (6)$$

In (5) and (6), only nearest-neighbor intersite couplings are considered. J_1 and J_2 are thus the two *different* nearest-neighbor Josephson couplings responsible for condensate tunneling. In principle, w_1, w_2 can be solved numerically, which in turn solve J_1, J_2 , and U . Typical value of J_1 or J_2 is less than 0.1 in the TB limit.

In the following, as mentioned before, we will focus on the breathing and phonon modes. Similar to Martikainen and Stoof¹², a Gaussian ansatz

$$\Phi_i(x, y, \ell; t) = \sqrt{\frac{N[1 + \delta_{\ell,i}(t)] B_0 [1 + \epsilon'_{\ell,i}(t)]}{\pi}} \\ \times \exp \left[-\frac{B_0 [1 + \epsilon'_{\ell,i}(t) + i\epsilon''_{\ell,i}(t)] (x^2 + y^2)}{2} + i\nu_{\ell,i}(t) \right] \quad (7)$$

is assumed for the wave functions, where N represents the average number of atoms per site and B_0 represents the condensate size in the equilibrium state. In fact, when N is fixed, the value of B_0 can be calculated through minimizing the GP energy in (5). This was done in Ref.¹² where $B_0 = 1/\sqrt{1 + 2U'}$ with $U' \equiv 4NU/\pi$ is given. No cross (xy) term is considered in (7) because breathing mode is isotropic in the x - y plane. For each site (ℓ, i) , dimensionless dynamical variable $\epsilon'_{\ell,i}(t)$, $\epsilon''_{\ell,i}(t)$, $\delta_{\ell,i}(t)$, and $\nu_{\ell,i}(t)$ corresponds respectively to fluctuations of the *local amplitude*, the *local phase*, the *number* of atoms, and the *global phase* of the condensate.

One crucial aspect on (7) is however that it enables naturally the coupling between the transverse breathing and longitudinal phonon modes¹². In a single Bose-condensed system with repulsive interaction, the fluctuation of condensed atom number is coupled to the

fluctuation of condensate size. When condensates are distributed in an optical lattice, the fluctuating degrees of freedom are complicated by the intersite coupling of Josephson tunnelling. For the present optical superlattice, the case is further complicated by the possible out-of-phase motion in addition to usual in-phase one. How important are the various couplings and how the condensates are fluctuating in such a system are thus the subject of the following studies.

By variational approach, one starts from the Lagrangian of the system, $L = T - E$. Here $T = \int d\mathbf{r} \frac{i\hbar}{2} (\Psi^* \partial \Psi / \partial t - \Psi \partial \Psi^* / \partial t)$ and E is given by (3) [and hence (5)]. Applying (7) in T and E and expanding to second order in dynamical variables, one obtains

$$\frac{T}{N} = \sum_{\ell} \sum_{i=1,2} \left\{ (1 + \delta_{\ell,i}) \left[-\dot{\nu}_{\ell,i} + \frac{\dot{\epsilon}_{\ell,i}''}{2} (1 - \epsilon'_{\ell,i}) \right] \right\}, \quad (8)$$

and

$$\frac{E}{N} = \sum_{\ell} \sum_{i=1,2} \left\{ (1 + \delta_{\ell,i}) \left[\frac{1}{2B_0} (2 + (\epsilon'_{\ell,i})^2) + \frac{B_0}{2} (\epsilon_{\ell,i}'')^2 + U' B_0 (1 + \epsilon'_{\ell,i}) \delta_{\ell,i} - \bar{\mu} \right] \right\} + E_J, \quad (9)$$

where the Josephson-tunneling term

$$\begin{aligned} E_J = & \frac{J_1}{4} \sum_{\ell} [(\epsilon'_{\ell,1} - \epsilon'_{\ell-1,2})^2 + 2(\epsilon_{\ell,1}'' - \epsilon_{\ell-1,2}'')^2 \\ & + 4(\nu_{\ell,1} - \nu_{\ell-1,2})^2 + (\delta_{\ell,1} - \delta_{\ell-1,2})^2 \\ & - 4(\nu_{\ell,1} - \nu_{\ell-1,2})(\epsilon_{\ell,1}'' - \epsilon_{\ell-1,2}'') - 4(\delta_{\ell,1} + \delta_{\ell-1,2})] \\ & + \frac{J_2}{4} \sum_{\ell} [(\epsilon'_{\ell,1} - \epsilon'_{\ell,2})^2 + 2(\epsilon_{\ell,1}'' - \epsilon_{\ell,2}'')^2 \\ & + 4(\nu_{\ell,1} - \nu_{\ell,2})^2 + (\delta_{\ell,1} - \delta_{\ell,2})^2 \\ & - 4(\nu_{\ell,1} - \nu_{\ell,2})(\epsilon_{\ell,1}'' - \epsilon_{\ell,2}'') - 4(\delta_{\ell,1} + \delta_{\ell,2})]. \end{aligned} \quad (10)$$

It is noted in (9) and (10) that when chemical potential $\bar{\mu}$ is chosen to be $\bar{\mu} = 3/2B_0 - B_0/2 - J_1 - J_2$ (as given by the stability condition of $\partial E / \partial \delta_{\ell,i} |_{\text{var.}=0} = 0$), linear in $\delta_{\ell,i}$ terms vanish. In this case, only second order terms are left with E_J in (10) and it justifies the smallness of E_J in accordance with TB.

With (8)-(10), one can then derive the Euler-Lagrange equations of motion for all eight dynamical variables. They can be linearized and written as

$$\begin{aligned} \dot{\epsilon}'_{\ell,1} &= 2B_0\epsilon_{\ell,1}'' + J_1(\epsilon_{\ell,1}'' - \epsilon_{\ell-1,2}'') + J_2(\epsilon_{\ell,1}'' - \epsilon_{\ell,2}''), \\ \dot{\epsilon}'_{\ell,2} &= -2B_0\epsilon_{\ell,2}'' + J_1(\epsilon_{\ell,2}'' - \epsilon_{\ell+1,1}'') + J_2(\epsilon_{\ell,2}'' - \epsilon_{\ell,1}''), \\ \dot{\epsilon}_{\ell,1}'' &= -\frac{2}{B_0}\epsilon'_{\ell,1} - 2U'B_0\delta_{\ell,1} - J_1(\epsilon'_{\ell,1} - \epsilon'_{\ell-1,2}) \\ &\quad - J_2(\epsilon'_{\ell,1} - \epsilon'_{\ell,2}), \\ \dot{\epsilon}_{\ell,2}'' &= -\frac{2}{B_0}\epsilon'_{\ell,2} - 2U'B_0\delta_{\ell,2} - J_1(\epsilon'_{\ell,2} - \epsilon'_{\ell+1,1}) \end{aligned}$$

$$\begin{aligned} & -J_2(\epsilon'_{\ell,2} - \epsilon'_{\ell,1}), \\ \dot{\delta}_{\ell,1} &= J_1[2(\nu_{\ell,1} - \nu_{\ell-1,2}) - (\epsilon_{\ell,1}'' - \epsilon_{\ell-1,2}'')] \\ & \quad + J_2[2(\nu_{\ell,1} - \nu_{\ell,2}) - (\epsilon_{\ell,1}'' - \epsilon_{\ell,2}'')], \\ \dot{\delta}_{\ell,2} &= J_1[2(\nu_{\ell,2} - \nu_{\ell+1,1}) - (\epsilon_{\ell,2}'' - \epsilon_{\ell+1,1}'')] \\ & \quad + J_2[2(\nu_{\ell,2} - \nu_{\ell,1}) - (\epsilon_{\ell,2}'' - \epsilon_{\ell,1}'')], \\ \dot{\nu}_{\ell,1} &= -(1 + 3U')B_0\epsilon'_{\ell,1} - 3B_0U'\delta_{\ell,1} \\ & \quad - \frac{J_1}{2}[(\delta_{\ell,1} - \delta_{\ell-1,2}) + (\epsilon'_{\ell,1} - \epsilon'_{\ell-1,2})] \\ & \quad - \frac{J_2}{2}[(\delta_{\ell,1} - \delta_{\ell,2}) + (\epsilon'_{\ell,1} - \epsilon'_{\ell,2})], \\ \dot{\nu}_{\ell,2} &= -(1 + 3U')B_0\epsilon'_{\ell,2} - 3B_0U'\delta_{\ell,2} \\ & \quad - \frac{J_1}{2}[(\delta_{\ell,2} - \delta_{\ell+1,1}) + (\epsilon'_{\ell,2} - \epsilon'_{\ell+1,1})] \\ & \quad - \frac{J_2}{2}[(\delta_{\ell,2} - \delta_{\ell,1}) + (\epsilon'_{\ell,2} - \epsilon'_{\ell,1})]. \end{aligned} \quad (11)$$




FIG. 2: Dispersions of four-branch modes resulting from (11). The parameters $J_1 = 0.09$ and $J_2 = 0.1$ are the same for all three frames, while $U' = 1, 10$, and 100 respectively for frame (a), (b), and (c). Here IB, OB, AP, and OP denote for in-phase breathing, out-of-phase breathing, acoustic phonon, and optical phonon modes. See text for more description.

Note that the above eight linear first-order differential equations can be cast into four second-order ones in terms of any two sets of variables $\epsilon'_{\ell,i}(t)$, $\epsilon_{\ell,i}''(t)$, $\delta_{\ell,i}(t)$, and $\nu_{\ell,i}(t)$ ($i = 1, 2$) only. (That is in the lowest-order limit, the other two sets will share the same excitation spectra.) Considering the coupled equations between $\epsilon'_{\ell,i}(t)$ and $\delta_{\ell,i}(t)$, for example, and searching for solutions of the type: $\epsilon'_{\ell,i}(t) \equiv \epsilon'_i e^{i(\ell kc - \omega t)}$ and $\delta_{\ell,i}(t) \equiv \delta_i e^{i(\ell kc - \omega t)}$, a 4×4 dynamical matrix (not shown) will be attained, which is then diagonalized to obtain four branches of modes. In Fig. 2, dispersions of the four branches of modes are shown for three cases: with same $J_1 = 0.09$ and $J_2 = 0.1$, but different $U' = 1, 10$, and 100 . In the study of Bose-Hubbard model on the SF-Mott insulator quantum phase transition in 3D, it has been established that $U/J = z \times 5.8$ (with z the number of nearest-neighbor sites) is a critical point for the case of one atom per site on average ($N = 1$). Below (above) it the system

is in the SF (insulating) phase¹⁴. Here we are interested in collective excitation of condensates with average atom number per site being $N \sim 10 - 100$. Thus the system being in the SF phase is justified when $U'/J < 100$ (recall $U' \sim NU$). Consequently the $U' = 1$ and 10 cases in Fig. 2 are considered to be in the SF phase, while the $U' = 100$ case may be close to the phase boundary or in the insulating phase.

The four branches of modes are labelled by in-phase breathing (IB), out-of-phase breathing (OB), acoustic phonon (AP), and optical phonon (OP) modes respectively (see Fig. 2). Strictly speaking, these descriptions are exact only when phonon modes are completely decoupled from breathing modes and at the same time, the propagating wave comprised of $\epsilon'_{\ell,1}(t)$, for instance, has a phase angle π shift (out-of-phase) by the the propagating wave comprised of $\epsilon'_{\ell,2}(t)$. The coupling effect, which is proportional to the magnitude of tunnelling strength J , is indeed small in the TB limit. One should be noted that the coupling effect is also accompanied by the factor of $B_0 U'$ (recall that B_0 corresponds to the size of condensate), to which it becomes more and more important in the large U' limit.

At the long-wavelength limit ($kc \ll 1$), dispersions of the four branches of modes can be analytically solved and uniformly written as

$$\omega_i^2 = \omega_{0i}^2 + u_i^2 k^2, \quad (12)$$

where, at the TB limit (keeping to J linear order), one obtains

$$\omega_{0i}^2 = \begin{cases} 4 + 8B_0(J_1 + J_2) \left(1 + U' + \frac{B_0^2 U'^2}{2}\right) & i = \text{OB} \\ 4 & i = \text{IB} \\ 8B_0 U'(J_1 + J_2) \left(1 - \frac{B_0^2 U'}{2}\right) & i = \text{OP} \\ 0 & i = \text{AP}, \end{cases} \quad (13)$$

and

$$u_{\text{IB}}^2 = -u_{\text{OB}}^2 = 2B_0(1 + U')c^2 \frac{J_1 J_2}{J_1 + J_2}$$

$$u_{\text{AP}}^2 = -u_{\text{OP}}^2 = 2B_0 U' c^2 \frac{J_1 J_2}{J_1 + J_2}. \quad (14)$$

As shown in (12)–(14), the slopes (u_i) and the $k = 0$ intercept (ω_{0i}) of the four branches of modes depend crucially on the values of J_1 , J_2 , and U' . Therefore observation of these modes will be a direct way to access them. Some features worth noting are as follows. For IB mode, the $k = 0$ frequency ($\omega = 2\omega_\rho$) is robust regardless of the values of J_1 and J_2 . It is a clear result to the uniform (single-condensate) limit. For AP mode, on the other hand, the phonon velocity at small k behaves like $u \sim \sqrt{B_0 U' J} c$, an expected result for the current system of a bulk modulus $\sim B_0 U' J$. In the large U' case [see Fig. 2(c)], IB and OP modes start to hybridize, to which a small gap (not visible in the scale potted) develops where the two branches cross. This feature may be a signature when the system undergoes a quantum phase transition from SF into insulating phase.

In summary, oscillations of Bose condensates in a 1D optical lattice with a two-point basis is investigated. Focuses are made to the transverse breathing modes and longitudinal phonon modes. Pertaining to condensates in site 1 and site 2, in-phase and out-of-phase modes are obtained. The mode dispersion relations depend crucially on the sizes of the two intersite Josephson tunneling strengths J_1 and J_2 and the on-site repulsion U . For the phonons, there are optical as well as acoustic modes, in a close resemblance to those in a 1D crystalline solid. While direct observation of these in-phase and out-of-phase oscillations of condensates may be resolution limited by current instruments, dispersion relations of these modes should be probable by inelastic scattering measurements.

We are grateful to Prof. D.-J. Han for stimulating discussion. This work is supported by National Science Council of Taiwan under Grant No. 94-2112-M-003-011.

¹ D. Jaksch, C. Bruder, J. Cirac, C.W. Gardiner, and P. Zoller, Phys. Rev. Lett. **81**, 3108 (1998).

² M. Greiner, O. Mandel, T. Esslinger, T.W. Hänsch, and I. Bloch, Nature **415**, 39 (2002).

³ M. Machholm, C.J. Pethick, and H. Smith, Phys. Rev. A **67**, 053613 (2003).

⁴ B. Wu and Q. Niu, Phys. Rev. A **61**, 023402 (2000).

⁵ J.I. Cirac and P. Zoller, Science **301**, 176 (2003).

⁶ K.G.H. Vollbrecht, E. Solano, and J. I. Cirac, Phys. Rev. Lett. **93**, 220502 (2004).

⁷ See, for example, F. Dalfovo, S. Giorgini, L.P. Pitaevskii,

and S. Stringari, Rev. Mod. Phys. **71**, 463 (1999).

⁸ M. Krämer, L. Pitavskii, and S. Stringari, Phys. Rev. Lett. **88**, 180404 (2002), and references therein.

⁹ K. Berg-Sørensen and K. Mølmer, Phys. Rev. A **58**, 1480 (1998).

¹⁰ J. Javanainen, Phys. Rev. A **60**, 4902 (1999).

¹¹ E. Taylor and E. Zaremba, Phys. Rev. A **68**, 053611 (2003).

¹² J.-P. Martikainen and H. T. C. Stoof, Phys. Rev. A **69**, 23608 (2004).

¹³ C. Schori, T. Stöferle, H. Moritz, M. Köhl, and T.

Esslinger, Phys. Rev. Lett. **93**, 240402 (2004).

H. T. C. Stoof, Phys. Rev. A **63**, 053601 (2001).

¹⁴ See, for example, D. van Oosten, P. van der Straten, and